



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[\[Sign In\]](#) [\[Regis\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Limits Preview/Index History Clipboard Details

About Entrez

All: 1 Review: 0

Text Version

☐ 1: [Eur J Pharmacol.](#) 2000 Sep 8;403(3):181-8.

[Related Articles, Links](#)

Entrez PubMed

Overview
Help | FAQ
Tutorials

New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources

Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

ELSEVIER
FULL-TEXT ARTICLE

Taurine attenuates hypertrophy induced by angiotensin II in cultured neonatal rat cardiac myocytes.

Azuma M, Takahashi K, Fukuda T, Ohyabu Y, Yamamoto I, Kim S, Iwao H, Schaffer SW, Azuma J.

Department of Clinical Evaluation of Medicines and Therapeutics, Graduate School of Pharmaceutical Sciences, Osaka University, Osaka 565,-0871, Suita, Japan.

The effect of taurine on angiotensin II-induced changes in cell morphology and biochemistry of the cultured neonatal cardiomyocyte was examined. Angiotensin II (1-100 nM) alone caused a slow increase in the surface area of the myocyte accompanied by an induction of the expression of atrial natriuretic peptide (ANP) and an upregulation of transforming growth factor beta(1) gene (TGF-beta(1)). The signaling pathway of angiotensin II (1-100 nM) was found to proceed through protein kinase C and the rapid activation of mitogen-activated protein (MAP) kinases. Pretreatment of the myocyte with taurine (20 mM) in the absence of angiotensin II had no visible effect on cell size or growth rate. However, the cells that were pretreated with taurine (20 mM) for 24 h exhibited reduced responsiveness to angiotensin II (100 nM) relative to surface cell area enlargement and the upregulation of the late and growth factor genes(ANP, TGF-beta(1)). Angiotensin II-mediated activation of the MAP kinases (extracellular signal-regulated protein kinase 1/2: ERK1/2) was not blocked by taurine. Taurine reduced the phosphorylation of a 29-kDa protein, a reaction which was enhanced by angiotensin II and appears to involve protein kinase C step. The results indicate that taurine is an effective inhibitor of certain aspects of angiotensin II action.

PMID: 10973617 [PubMed - indexed for MEDLINE]

Jun 21 2006 12:14:26